

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Chelating and Bridging Roles of 2-(2-Pyridyl)benzimidazole and Bis(diphenylphosphino)acetylene in Stabilizing a Cyclic Tetranuclear Platinum(II) Complex.

Permalink

<https://escholarship.org/uc/item/9n91c5mj>

Journal

Inorganic chemistry, 58(21)

ISSN

0020-1669

Authors

Nabavizadeh, S Masoud
Niroomand Hosseini, Fatemeh
Niknam, Fatemeh
et al.

Publication Date

2019-11-01

DOI

10.1021/acs.inorgchem.9b02274

Supplemental Material

<https://escholarship.org/uc/item/9n91c5mj#supplemental>

Peer reviewed

Chelating and Bridging Roles of 2-(2-Pyridyl)-benzimidazole and Bis(diphenylphosphino)acetylene in Stabilizing a Cyclic Tetranuclear Platinum(II) Complex

S. Masoud Nabavizadeh,^{*a,b} Fatemeh Niroomand Hosseini,^c Fatemeh Niknam,^a Peyman Hamidizadeh,^a Peter C Ford^b and Mahdi M. Abu-Omar^{*b}

^aProfessor Rashidi Laboratory of Organometallic Chemistry, Department of Chemistry, College of Sciences, Shiraz University, Shiraz, 71467-13565, Iran, ^bDepartment of Chemistry and Biochemistry, University of California, Santa Barbara, Santa Barbara, California 93106, United States. ^cDepartment of Chemistry, Shiraz Branch, Islamic Azad University, Shiraz 71993–37635, Iran

Abstract

The reaction of complex [Pt(Me)(DMSO)(pbz)], **1**, (pbz = 2-(2-pyridyl)benzimidazolate) with [PtMe(Cl)(DMSO)₂], **B**, followed by addition of bis(diphenylphosphino)acetylene (dppac), gave the novel tetranuclear platinum complex [Pt₄Me₄(μ-dppac)₂(pbz)₂Cl₂], **2**, bearing both the pbz and dppac ligands. In this structure, the pbz ligands act as both chelating and bridging to stabilize the tetraplatinum framework. The tetranuclear Pt(II) complex was fully characterized by NMR spectroscopy, X-ray crystallography and mass spectrometry and its electronic structure was investigated and supported by DFT calculations.

Introduction

The construction of multinuclear metal assemblies has been a growing area of research during the past decade.¹ Developments based on the use of different ligands as the building unit for the synthesis of compounds have been reported for binuclear Pt complexes,² although to the of our knowledge, tetranuclear complexes have been rarely prepared.³

Wilson and Lippard reported⁴ a tetranuclear platinum complex in which two acetate-bridged diplatinum(III) units are connected by two bridging amido ligands (**I**, Chart 1). The syntheses of tetranuclear platinum complexes in which the metal centers are supported by phosphido ligands have been reported by Forniés and coworkers (**II**, Chart 1).⁵ Stang and Cao reported⁶ the preparation and characterization, via self-assembly, of platinum- and palladium-based cationic, tetranuclear, macrocyclic complexes (**III**, Chart 1). In these complexes, 4,4'-bipyridine acts as spacer ligand between Pt centers. Youngs and his coworkers also reported⁷ the synthesis of tetraplatinum square heterocycle which had large cavities and in which the rings were composed of only platinum and butadiyne fragments (**IV**, Chart 1). They used chelating phosphines in order to enforce the required *cis* geometry at the platinum atom. Yam et. al.⁸ synthesized a V-shape luminescent tetranuclear platinum(II) alkynyl complex (**V**, Chart 1) and its electrochemical behavior was studied by cyclic voltammetry. Also, very recently, a tetranuclear Pt(II) compound using the 1,2,3-tetrazolate as a building block (**VI**, Chart 1) is reported⁹ in which all triazolate nitrogen atoms are coordinated to 3 different Pt(II) ions. Casas and his coworker synthesized¹⁰ a tetranuclear platinum(II)-platinum(IV) mixed-valence complexes by addition of the stoichiometric amount of a Pt(II) reagent [*cis*-Pt(C₆F₅)₂(thf)₂] to some binuclear platinum(IV) complexes (**VII**, Chart 1).




Chart 1. Examples of some tetraplatinum complexes

On the other hand, cyclometalated complexes have attracted attention because of their several applications in science.¹¹ Of particular interest are cycloplatinated complexes bearing strong field ligands such as benzo[h]quinolate (bhq) and 2-phenylpyridinate (ppy).¹² The

cycloplatinated(II) compounds with the $[\text{Pt}(\text{C}^{\wedge}\text{N})\text{R}]$ moiety ($\text{C}^{\wedge}\text{N}$ = cyclometalated ligand and R = alkyl, aryl or halide) are very attractive due to their ease of preparation through C-H activation of cyclometalated ligands like Hppy or Hbhq with a suitable Pt(II) precursor complex preparation.^{12c-f, 13} Earlier we studied the reaction of the double rollover cycloplatinated(II) complex $[\text{Pt}_2(\mu\text{-bpy-2H})(\text{Me})_2(\kappa^1\text{-dppm})_2]$, bpy= 2,2'-bipyridine, (in which each metallic center bears a monodentate dppm ligand) with the known monomeric cycloplatinated (II) complex $[\text{Pt}(\text{Me})(\text{bhq})(\text{SMe}_2)]$ in a 1:2 molar ratio, which results in the unsymmetrical tetranuclear complex $[\{(\text{bhq})(\text{Me})\text{Pt}(\mu\text{-dppm})\text{Pt}(\text{Me})\}_2(\mu\text{-bpy-2H})]$ (**VIII**, Chart 1).^{13b}

The 2-(2-pyridyl)benzimidazole (Hpbz) ligand normally acts as an anionic $\text{N}^{\wedge}\text{N}$ bidentate ligand through N-H bond activation of the imidazole unit (instead of C-H activation in Hppy and Hbhq ligands).¹⁴ Hpbz derivatives have been also used as 'bipyridine analogues' in some complexes with metal centers such as Cu(I,II),¹⁵ Zn(II),¹⁶ Ru(II),¹⁷ Os(II),¹⁸ and Re(I).¹⁹ The ease of synthesis and the commercial availability of Hpbz are the factors that make this an attractive ligand. Although there are many reports on preparation of metal complexes in which the pbz acts as $\text{N}^{\wedge}\text{N}$ bidentate ligand, the complexes bearing this ligand as a bridging ligand between two metal centers have not been explored.

Although the ability of the biphosphine ligands to form polynuclear metal complexes, in which the metal centers are supported by these bridging ligands, has produced a very rich and interesting chemistry,^{13c, 20} best of our knowledge, there is no report on using 2-(2-pyridyl)-benzimidazole where it can concurrently act as a chelating and bridging ligand. These, together with our current interest in the design and synthesis of organoplatinum(II) complexes²¹ have prompted us to exploit the 2-(2-pyridyl)-benzimidazole and bis(diphenylphosphino)acetylene (dppac) units as the backbone for the construction of a cyclic tetraplatinum(II) complexes. In fact, an attempt has been made to synthesize an interesting tetranuclear organoplatinum(II)

complex in which, for the first time, pbz ligands act as both chelating and bridging to stabilize the tetraplatinum framework. In this complex dppac ligands are also located as bridging ligands between Pt(II) centers. The electronic structure of this complex, supported by theoretical calculations, is also investigated.

Experimental section

General remarks

The ^1H and ^{31}P NMR spectra were recorded in CDCl_3 solutions on a Bruker Avance DPX 400 MHz spectrometer using TMS (^1H) and H_3PO_4 (^{31}P) as references. All chemical shifts and coupling constants are in ppm and Hz, respectively. Microanalyses were performed using a Thermo Finnigan Flash EA-1112 CHNSO rapid elemental analyzer and melting points were recorded on a Buchi 530 apparatus. UV-vis absorption spectra were recorded on a PerkinElmer Lambda 25 spectrophotometer using a cuvette with a path length of 1 cm and/or 1 mm. The complexes $[\text{Pt}(\text{Me})_2(\text{DMSO})_2]$, **A**,²² and $[\text{Pt}(\text{Me})(\text{DMSO})(\text{pbz})]$, **1**,²³ were prepared by the literature method. Bis(diphenylphosphino)acetylene (dppac) and 2-(2-pyridyl) benzimidazole (Hpbz) were purchased from commercial sources.

Synthesis of Pt complexes

$[\text{Pt}(\text{Me})\text{Cl}(\text{DMSO})_2]$, **B**

This compound had been previously synthesized by the reaction of $[\text{PtCl}_2(\text{DMSO})_2]$ with SnMe_4 for 24 h with a yield of 55%.²⁴ In an alternative method, 381 mg (1.00 mmol) of **A** and 421 mg (1.00 mmol) of $[\text{PtCl}_2(\text{DMSO})_2]$ were added to 50 mL chloroform or acetone and stirred for 6 h. The solvent was then removed by evaporation to give 640 mg **B** (1.59 mmol). Yield: 80.0%; ^1H NMR in CDCl_3 ; *trans* isomer: δ 0.99 [s, $^2J_{\text{PtH}} = 78.9$ Hz, 3H, Me]; 3.42, [s, $^3J_{\text{PtH}} = 24.23$ Hz, 12H, Me of DMSO]; *cis* isomer: δ 1.17 [s, $^2J_{\text{PtH}} = 79.0$ Hz, 3H, Me]; 3.21, [s, $^3J_{\text{PtH}} = 8.8$ Hz, 6H, DMSO *trans* to Me]; 3.49 [s, $^3J_{\text{PtH}} = 38.2$ Hz, 6H, DMSO *trans* to Cl].

[Pt₄(Me)₄(μ-dppac)₂(pbz)₂Cl₂], **2**




Chart 2. NMR labeling for complex **2**

[Pt(Me)Cl(DMSO)₂], **B**, (401 mg, 1.00 mmol) was added to a solution of **1**, (482 mg, 1.00 mmol) in acetone (100 ml). This was stirred at room temperature for 24 h and then dppac (394 mg, 1 mmol) was added to the solution. After 20 h solvent was removed and the yellow residue was washed with cold diethylether to give 800 mg **2** (0.381 mmol). Yield: 76.0%; m.p. > 250 °C. Anal. Calc. for C₈₀H₆₈Cl₂N₆Pt₄P₄: C, 46.0; H, 3.2; N, 4.0 %, Found, C, 45.8; H, 3.4; N, 4.1%, NMR in CDCl₃: δ(¹H) 0.56 [d, ²J_{PH} = 83.5 Hz, ³J_{PH} = 7.3 Hz, 6H, Me ligands *trans* to N of 2-pyridyl ring], 1.09 [d, ²J_{PH} = 72.6 Hz, ³J_{PH} = 4.7 Hz, 6H, Me ligands *trans* to N of benzimidazole ring]; 8.15 [dd, ³J_{PH} = 43.6 Hz, ³J_{HH} = 6.5 Hz, ⁴J_{PH} = 5.3 Hz, 2H, H⁶]; 7.84 and 7.33 [m, Ph of dppac]; 7.45 [m, H^{6'} and H^{3'}]; 6.90 [d, ³J_{HH} = 5.4 Hz, 2H, H³]; 6.79 [t, ³J_{HH} = 15.2 Hz, 2H, H^{5'}]; 6.58 [t, ³J_{HH} = 12.7 Hz, 2H, H⁵]; 6.31 [t, ³J_{HH} = 15.1 Hz, 2H, H^{4'}]; 6.07 [d, ³J_(HH) = 8.6 Hz, 2H, H⁴]. δ(³¹P) 27.9 [s, ¹J_{PTP} = 2545 Hz, 2P, *trans* to benzimidazole ring], 20.6 [s, ¹J_{PTP} = 4392 Hz, 2P, *trans* to Cl]. ESI-TOF-MS: m/z 2107.4 [**2** + H₃O]⁺, 1717.4 [**2** – dppac + Na]⁺.

Crystallographic data

Single crystal X-ray diffraction data for complex **2** were collected on a Bruker KAPPA APEX II diffractometer equipped with an APEX II CCD detector using a TRIUMPH

monochromator with a Mo K α X-ray source ($k = 0.71073 \text{ \AA}$). The crystal was mounted on a cryoloop under Paratone-N oil and kept under nitrogen. Absorption correction of the data was carried out using the multiscan method SADABS.²⁵ Subsequent calculations were carried out using SHELXTL.²⁶ Structure determination was done using intrinsic methods. Structure solution, refinement, and creation of publication data was performed using SHELXTL. Crystallographic information is presented in Table 1. Crystallographic data for the structural analysis has been deposited with the Cambridge Crystallographic Data Centre, No. CCDC 1874226. Copies of this information may be obtained free of charge from: The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK. Fax: +44(1223)336-033, e-mail: deposit@ccdc.cam.ac.uk, or www.ccdc.cam.ac.uk.

Table 1. Crystal and structure refinement for complex **2.4(CH₃)₂CO**.

Empirical formula	C ₉₂ H ₉₂ Cl ₂ N ₆ O ₄ P ₄ Pt ₄
Formula weight	2320.85
Temperature (K)	100(2)
Wavelength(Å)	0.71073
Crystal system	Triclinic
Space group	P-1
<i>a</i> /Å	11.207(3)
<i>b</i> /Å	11.658(3)
<i>c</i> /Å	17.085(5)
β /°	94.774(7)
VolumeÅ ³	2151.6(10)
<i>Z</i>	1
Density (calculated)/ Mg/m ³	1.791
Absorption coefficient/mm ⁻¹	6.672
F(000)	1124
Theta range for data collection	1.229 to 27.266°
Reflections collected	18876
Independent reflections, [R(int)]	9565, [0.0356]
Max. and min. transmission	0.7455 and 0.5607
Data / restraints / parameters	9565 / 0 / 511
Goodness-of-fit on F ²	0.976
R1/wR2 [I>2sigma(I)]	0.0294/ 0.0596


Computational details

Gaussian 09 was used²⁷ to fully optimize the structure at the B3LYP level of density functional theory. The effective core potential of Hay and Wadt with a double- ξ valence basis set (LANL2DZ) was chosen to describe Pt.²⁸ The 6-31G(d) basis set was used for all other atoms. To evaluate and ensure the optimized structures of the molecules, frequency calculations were carried out using analytical second derivatives. In all cases only real frequencies were obtained for the optimized structures. The calculations for the electronic absorption spectra by time-dependent DFT (TD-DFT) were performed at the same level of theory. The compositions of molecular orbitals and theoretical absorption spectra were plotted using the “Chemissian” software.²⁹

Results and Discussion

Complex synthesis and characterization

The detailed synthetic route for the tetranuclear complex $[\text{Pt}_4(\text{Me})_4(\mu\text{-dppac})_2(\text{pbz})_2\text{Cl}_2]$, **2**, is depicted in Scheme 1. The precursor complex $[\text{Pt}(\text{Me})(\text{DMSO})(\text{pbz})]$, **1**, was prepared through the reaction of $[\text{Pt}(\text{Me})_2(\text{DMSO})_2]$, **A**, with Hpbz in 1:1 ratio. In the key step complex **B** was added to a solution of **1** in acetone to produce an intermediate which is suggested to be the dimeric Pt complex $[\text{Cl}(\text{Me})(\text{DMSO})\text{Pt}(\mu\text{-pbz})\text{Pt}(\text{Me})(\text{DMSO})]$, **C**. Then, dppac was added in situ to the solution to result in the assembly of **2** as a yellow solid.



Scheme 1. Route to preparation of complex **2**

The structure of the new complex **2** was fully characterized using multinuclear NMR spectroscopy (^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR), electrospray ionization mass spectroscopy (ESI-MS) and X-ray crystallography. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum is illustrated in Fig. 1A (details of the numerical NMR data are collected in the Experimental section). The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **2** displays two distinct singlets with equal intensities, having platinum satellites. They appear at $\delta = 20.6$ ($^1J_{\text{PtP}} = 4392$ Hz) and 27.9 ($^1J_{\text{PtP}} = 2545$ Hz), which are respectively assigned to the P *trans* to Cl and the P *trans* to ligating N atom of benzimidazole ring, in agreement with high *trans* influence of N compared to Cl. It should be noted that the $^1J_{\text{PtP}}$ values of 4264 and 3081 Hz were previously reported for P atom *trans* to Cl in complexes $[\text{Pt}(\text{Me})(\text{dppe})\text{Cl}]$ (dppe = $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2$) and $[\text{PtCl}_2(\text{dppm})]$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_2$),³⁰ respectively. On the basis of the ^1H NMR spectrum of **2** (see Fig. 1B), two different Me groups are observed, being *trans* to

pyridyl and benzimidazole rings. The Me group *trans* to ligating N of pyridyl was observed to be appeared at $\delta = 0.56$ as a doublet signal with $^2J_{\text{PtH}} = 83.5$ Hz and $^3J_{\text{PH}} = 7.3$ Hz. Whilst, the Me group *trans* to ligating N of benzimidazole ring was appeared at $\delta = 1.09$ ppm with a considerably lower $^2J_{\text{PtH}} = 72.6$ Hz and $^3J_{\text{PH}} = 4.7$ Hz values. It indicates more *trans* influence of ligating N of benzimidazole ring in comparison to that for 2-pyridyl ring.




Fig. 1 (A) ^{31}P and (B) ^1H (aliphatic region) NMR spectra of the complex **2**.

Crystal Structure

Yellow crystals of complex **2** were obtained upon diffusion of *n*-hexane vapor into the acetone solution. The structure obtained by X-ray crystallography technique is presented in Fig. 2 and the numerical data are summarized in Table 1 (see the caption of Fig. 2 for selected geometrical parameters). To the best of our knowledge, the complex **2** is the first example of a tetranuclear Pt complex having both bridging 2-(2'-pyridyl)-benzimidazolate and bis(diphenylphosphino)acetylene ligands. In this structure, pbz ligands act as both chelating and bridging ligand to stabilize the tetraplatinum framework. The molecular structure of complex **2** certifies that the $\text{Pt}_4(\text{dppac})_2(\text{pbz})_2$ framework was formed. As shown in Fig. 2, the solid state structure exhibits distorted four-coordinated square planar geometry around the platinum centers. The methyl groups are *trans* to the pyridyl and benzimidazole rings of pbz

ligand. The largest deviation from ideal square planar geometry belongs to the N3-Pt2-N2 angle at $78.01(13)^\circ$ which is associated with the Pt(pbz) chelate ring. One of P atoms of dppac ligand is located *trans* to ligating N of benzimidazole ring (Pt2-P2 = 2.1910(13) Å) and another one is *trans* to Cl atom (Pt1-P1 = 2.1754(12) Å). The Me groups are located *trans* to the ligating N of the pyridyl (Pt2-C26 = 2.054(4) Å) and benzimidazole (Pt1-N1 = 2.122(3) Å) rings (See Fig. 2). Therefore the structure of complex **2** confirms that “Pt(μ -pbz)Pt” moieties are connected through dppac bridging ligand in which the P-C-C-P fragment is almost linear. The terminal Me ligands are *transoid* in relation to each other. It should be noted that this kind of bridged dppac complexes where a dppac molecule acts as a bridging ligand between two metal fragments are not common.^{2a, 30-31}




Fig. 2 (Top) Structure of complex **2**. All hydrogen atoms and solvent molecules are omitted for clarity. Selected geometrical parameters (Å, deg.): Pt1-Cl1 2.3597(12); Pt1-C1 2.064(4);

Pt1-P1 2.1754(12); Pt1-N1 2.122(3); Pt2-N2 2.078(4); Pt2-N3 2.134(3); Pt2-P2 2.1910(13); Pt2-C26 2.054(4); Cl1-Pt1-N1 88.28(10); Cl1-Pt1-P1 175.11(4); Cl1-Pt1-C1 86.89(12); N1-Pt1-P1 96.52(10); P2-Pt2-N3 104.73(10); P2-Pt2-C26 83.69(13); N3-Pt2-N2 78.01(13); N2-Pt2-C26 93.63(16). (Bottom) 18-membered Pt₄P₄N₄C₆ central core of tetranuclear complex **2**.

ESI-MS result further confirms the molecular formulae formation by demonstrated isotopically well-resolved peak at $m/z = 2107.4$ which is related to $[\mathbf{2} + \text{H}_3\text{O}]^+$. The peak matched well with its calculated theoretical isotope distributions (see Fig. 3).




Fig. 3 Expanded (left) and simulated (right) isotopic pattern ESI-MS spectrum of complex $[\mathbf{2} + \text{H}_3\text{O}]^+$, showing the expected intensity due to isotopic distribution.

Absorption spectrum

Complex **2** is stable in most of common organic solvents including acetone, CH_2Cl_2 , CHCl_3 and CH_3CN which allows the investigation of its absorption spectrum in a wide range of solvents. Fig. 4A shows the absorption spectrum of **2** in different solvents; the numerical data are listed in Table 2. The complex **2** shows intense absorption bands in the 300-400 nm

region I all solvents. As an example, two absorption bands at 312 and 358 nm ($\epsilon = 3.00 \times 10^4$ and $2.60 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$, respectively) appear in CH_2Cl_2 solvent. In order to gain further insight into the nature of electronic transitions for the complex **2** in CH_2Cl_2 , TD-DFT calculations were performed on the optimized ground state geometries. As shown in Fig. 4B, the computed absorption wavelengths (bars) are in good agreement with its experimental spectrum. The numerical data for the selected calculated wavelengths together with their assignments are listed in Table 2. For the complex **2**, the plots of molecular orbitals, the energies of the selected molecular orbitals together with their compositions in terms of ligands and metals are shown in Fig 5. The absorption bands at higher energies are attributed to L'LCT, IL and MLCT (L=pbz and L'= Cl).




Fig. 4 (A) UV-visible absorption spectrum of complex **2** at different solvents ($1.0 \times 10^{-5} \text{ M}$); dash line shows the UV-visible spectrum of acetone, UV cutoff of acetone is 330 nm. (B) Overlaid experimental absorbance (spectrum) and calculated TD-DFT (bar) for the complex **2** in CH_2Cl_2 .

Table 2. TD-DFT computed spectra and experimental wavelength (nm) for the complex **2** (1.0×10^{-5} M) at the ground state geometry in CH_2Cl_2 solution (only transitions with a probability higher than 20% are reported).

$\lambda_{\text{cal}} (f)$	$\lambda_{\text{max}}(\text{exp})$	Transitions (probability)	Assignments
345 (0.480)	312	HOMO-3 – LUMO+1 (46)	(L=pbz, L'= Cl, L'' = dppac) L'LCT, MLCT, IL
		HOMO-2 – LUMO (35)	IL, L'LCT, MLCT
365 (0.150)	358	HOMO – LUMO (63)	L'LCT, MLCT
		HOMO-1 – LUMO+1 (29)	L'LCT, MLCT




Fig. 5 Representative frontier orbitals and the relative composition energy levels for complex **2**.

The partial molecular orbital diagram for the tetranuclear platinum complex **2** with the several highest occupied and lowest unoccupied molecular orbital contours are depicted in Fig. 5 and the distribution of important frontier orbitals in the ground state and the corresponding partial

molecular orbital composition (percentages) are provided. The values of the HOMO–LUMO energy gap is equal to 3.588 eV. Analysis of the frontier molecular orbitals reveals that the maximum contribution in HOMO and HOMO-1 (the highest occupied molecular orbital) belongs to the Cl ligand and Pt metal center (in PtMeCINP moiety) with small contribution of dppac ligand. In addition, the results show that LUMO and LUMO+1 (lowest unoccupied molecular orbitals) are significantly localized on pbz bridging ligand, and LUMO+2 is contributed on dppac ligand.

Conclusions

Reaction of complex [Pt(Me)(DMSO)(pbz)], in which pbz = 2-(2-pyridyl)benzimidazolate, with [Pt(Me)Cl(DMSO)₂] in the presence of bis(diphenylphosphino)acetylene (= dppac) leads to a rare tetranuclear Pt(II) complex in which two 2-(2-pyridyl)-benzimidazolate ligands act as both chelating and bridging ligands to stabilize the tetraplatinum(II) complex. Dppac ligands are coordinated between Pt centers in a bridging mode *via* P-donor atoms, making a cyclic tetranuclear platinum complex. Investigation of electronic structure and related DFT and TD-DFT calculations supported a mixed MLCT (metal-to-ligand charge transfer) and L'LCT (ligand-to-ligand charge transfer) and IL (Intra ligand) transition (L=pbz, and L'= Cl).

ASSOCIATED CONTENT

Supporting Information

The computational details for complex **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*Email: nabavizadeh@shirazu.ac.ir (S.M. Nabavizadeh), abuomar@chem.ucsb.edu (M.M. Abu-Omar).

ORCID

S. Masoud Nabavizadeh: [0000-0003-3976-7869](https://orcid.org/0000-0003-3976-7869)

Fatemeh Niroomand Hosseini: [0000-0002-5856-8104](https://orcid.org/0000-0002-5856-8104)

Mahdi M. Abu-Omar: [0000-0002-4412-1985](https://orcid.org/0000-0002-4412-1985)

Peter C Ford: [0000-0002-5509-9912](https://orcid.org/0000-0002-5509-9912)

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interests.

Acknowledgments

We thank the Iran National Science Foundation (Grant No. 97012633), Shiraz University, the Islamic Azad University (Shiraz Branch) and Department of Chemistry and Biochemistry at UCSB for financial support.

References

1. (a) Holliday, B. J.; Mirkin, C. A., Strategies for the construction of supramolecular compounds through coordination chemistry. *Angew, Chem. Inter. Ed.* **2001**, *40*, 2022-2043; (b) Wu, G.-Y.; Chen, L.-J.; Xu, L.; Zhao, X.-L.; Yang, H.-B., Construction of supramolecular hexagonal metallacycles via coordination-driven self-assembly: Structure, properties and application. *Coord. Chem. Rev.* **2018**, *369*, 39-75.
2. (a) Nabavizadeh, S. M.; Sepehrpour, H.; Jamshidi, M.; Hosseini, F. N.; Owczarzak, A. M.; Kubicki, M., Binuclear organoplatinum(II) complexes with double bis(diphenylphosphino)acetylene

bridges: Synthesis, X-ray structure determination, electronic structures and DFT calculations. *J. Organomet. Chem.* **2016**, *808*, 34-41; (b) Qu, L.; Li, C.; Shen, G.; Gou, F.; Song, J.; Wang, M.; Xu, X.; Zhou, X.; Xiang, H., Syntheses, crystal structures, chirality and aggregation-induced phosphorescence of stacked binuclear platinum (II) complexes with bridging Salen ligands. *Mater. Chem. Front.* **2019**; (c) Hoseini, S. J.; Fath, R. H.; Fard, M. A.; Behnia, A.; Puddephatt, R. J., A Bridging Peroxide Complex of Platinum (IV). *Inorg. Chem.* **2018**, *57*, 8951-8955; (d) Niroomand Hosseini, F., Determination of Reaction Kinetic Parameters from Variable Temperature Kinetic Study for Oxidative Addition Reaction on Binuclear Cyclometalated Platinum (II) Complexes. *Inorg. Chem. Res.* **2019**, *2*, 26-31.

3. (a) Zheng, Q.; Schneider, J. F.; Amini, H.; Hampel, F.; Gladysz, J. A., Wire like diplatinum, triplatinum, and tetraplatinum complexes featuring X [PtC≡CC≡CC≡CC≡C]_mPtX segments; iterative syntheses and functionalization for measurements of single molecule properties. *Dalton Trans.* **2019**, 48, 5800-5816; (b) Owen, G. R.; Hampel, F.; Gladysz, J., Syntheses and structures of tetraplatinum bis (polyynediyl) complexes with laterally arrayed sp carbon chains. *Organometallics* **2004**, *23*, 5893-5895; (c) Yamaguchi, T.; Saito, H.; Maki, T.; Ito, T., Syntheses and Structures of Bridge and Chelate Isomers of Tetraplatinum(II) Cluster Complex with Diethyldithiophosphate Ion (Et₂dtP) at In-Plane Sites, [Pt₄(CH₃COO)₄(Et₂dtP)₄], and Their Mutual Isomerization via Cluster Core Rotation. *J. Am. Chem. Soc.* **1999**, *121*, 10738-10742.

4. Wilson, J. J.; Lippard, S. J., Acetate-bridged platinum (III) complexes derived from cisplatin. *Inorg. Chem.* **2012**, *51*, 9852-9864.

5. Forniés, J.; Fortuño, C.; Gil, R.; Martín, A., Tetranuclear platinum phosphido complexes with different structures. *Inorg. Chem.* **2005**, *44*, 9534-9541.

6. Stang, P. J.; Cao, D. H., Transition metal based cationic molecular boxes. Self-assembly of macrocyclic platinum (II) and palladium (II) tetranuclear complexes. *J. Am. Chem. Soc.* **1994**, *116*, 4981-4982.

7. ALQaisi, S. M.; Galat, K. J.; Chai, M.; Ray, D. G.; Rinaldi, P. L.; Tessier, C. A.; Youngs, W. J., Synthesis of Neutral Tetranuclear and Octanuclear Macrocyclic Platinum- Butadiyne Heterocyclines. *J. Am. Chem. Soc.* **1998**, *120*, 12149-12150.

8. Tao, C.-H.; Wong, K. M.-C.; Zhu, N.; Yam, V. W.-W., Versatile precursors for multinuclear platinum (ii) alkynyl assembly—synthesis, structural characterization and electrochemical studies of luminescent platinum(II) alkynyl complexes. *New J. Chem.* **2003**, *27*, 150-154.

9. Komeda, S.; In, Y.; Tomoo, K.; Minoura, K.; Sato, T.; Reedijk, J.; Ishida, T.; Chikuma, M., Associative intraligand substitution of anticancer azolato-bridged compounds without a square-pyramidal intermediate: formation of a unique tetranuclear, μ_3 -1, 2, 3-triazolato-N1, N2, N3-bridged Pt (II) compound. *Inorg. Chim. Acta* **2019**, *495*, 118999.

10. Casas, J. M.; Diosdado, B. E.; Forniés, J.; García-Monforte, M. Á.; Laporta, R.; Martín, A.; Tomás, M., Synthesis and characterization of binuclear Pt(IV) complexes and tetranuclear mixed valence complexes of Platinum(II)-Platinum(IV). *J. Organomet. Chem.* **2019**, *897*, 130-138.

11. (a) Li, K.; Tong, G. S. M.; Wan, Q.; Cheng, G.; Tong, W.-Y.; Ang, W.-H.; Kwong, W.-L.; Che, C.-M., Highly phosphorescent platinum (II) emitters: photophysics, materials and biological applications. *Chem. Sci.* **2016**, *7*, 1653-1673; (b) Lanoë, P.-H.; Moreno-Betancourt, A.; Wilson, L.; Philouze, C.; Monnereau, C.; Jamet, H.; Jouvenot, D.; Loiseau, F., Neutral heteroleptic cyclometallated Platinum (II) complexes featuring 2-phenylbenzimidazole ligand as bright emitters in solid state and in solution. *Dyes and Pigm.* **2019**, *162*, 967-977; (c) Zhao, Q.; Huang, C.; Li, F., Phosphorescent heavy-metal complexes for bioimaging. *Chem. Soc. Rev.* **2011**, *40*, 2508-2524; (d) Li, T.-Y.; Liang, X.; Zhou, L.; Wu, C.; Zhang, S.; Liu, X.; Lu, G.-Z.; Xue, L.-S.; Zheng, Y.-X.; Zuo, J.-L., N-heterocyclic carbenes: versatile second cyclometalated ligands for neutral iridium (III) heteroleptic complexes. *Inorg. Chem.* **2014**, *54*, 161-173; (e) Omae, I., Applications of five-membered ring products of cyclometalation reactions as anticancer agents. *Coord. Chem. Rev.* **2014**, *280*, 84-95.

12. (a) Mdeleleni, M. M.; Bridgewater, J. S.; Watts, R. J.; Ford, P. C., Synthesis, structure, and spectroscopic properties of ortho-metalated platinum (II) complexes. *Inorg. Chem.* **1995**, *34*, 2334-2342; (b) Fereidoonhezad, M.; Kaboudin, B.; Mirzaee, T.; Babadi Aghakhanpour, R.; Golbon

- Haghighi, M.; Faghih, Z.; Faghih, Z.; Ahmadipour, Z.; Notash, B.; Shahsavari, H. R., Cyclometalated Platinum (II) Complexes Bearing Bidentate O, O'-Di (alkyl) dithiophosphate Ligands: Photoluminescence and Cytotoxic Properties. *Organometallics* **2017**, *36*, 1707-1717; (c) Shahsavari, H. R.; Babadi Aghakhanpour, R.; Nikraves, M.; Ozdemir, J.; Golbon Haghighi, M.; Notash, B.; Beyzavi, M. H., Highly Emissive Cycloplatinated (II) Complexes Obtained by the Chloride Abstraction from the Complex [Pt(ppy)(PPh₃)(Cl)]: Employing Various Silver Salts. *Organometallics* **2018**, *37*, 2890-2900; (d) Sangari, M. S.; Haghighi, M. G.; Nabavizadeh, S. M.; Pfitzner, A.; Rashidi, M., Influence of ancillary ligands on the photophysical properties of cyclometalated organoplatinum (ii) complexes. *New J. Chem.* **2018**, *42*, 8661-8671; (e) Molaei, H.; Nabavizadeh, S. M.; Jamshidi, M.; Vilsmeier, M.; Pfitzner, A.; Sangari, M. S., Phosphorescent heterobimetallic complexes involving platinum (IV) and rhenium (VII) centers connected by an unsupported μ -oxido bridge. *Dalton Trans.* **2017**, *46*, 16077-16088; (f) Berenguer, J. R.; Lalinde, E.; Moreno, M. T., Luminescent cyclometalated-pentafluorophenyl Pt II, Pt IV and heteropolynuclear complexes. *Coord. Chem. Rev.* **2018**, *366*, 69-90.
13. (a) Millán, G.; Giménez, N.; Lara, R.; Berenguer, J. s. R.; Moreno, M. T.; Lalinde, E.; Alfaro-Arnedo, E.; López, I. a. P.; Piñero-Hermida, S.; Pichel, J. G., Luminescent Cycloplatinated Complexes with Biologically Relevant Phosphine Ligands: Optical and Cytotoxic Properties. *Inorg. Chem.* **2019**, *58*, 1657-1673; (b) Aghakhanpour, R. B.; Nabavizadeh, S. M.; Rashidi, M., Newly designed luminescent di-and tetra-nuclear double rollover cycloplatinated (II) complexes. *J. Organomet. Chem.* **2016**, *819*, 216-227; (c) Aghakhanpour, R. B.; Nabavizadeh, S. M.; Rashidi, M.; Kubicki, M., Luminescence properties of some monomeric and dimeric cycloplatinated (II) complexes containing biphosphine ligands. *Dalton Trans.* **2015**, *44*, 15829-15842; (d) Schneider, L.; Sivchik, V.; Chung, K.-y.; Chen, Y.-T.; Karttunen, A. J.; Chou, P.-T.; Koshevoy, I. O., Cyclometalated platinum (II) cyanometallates: luminescent blocks for coordination self-assembly. *Inorg. Chem.* **2017**, *56*, 4459-4467; (e) Barzegar-Kiadehi, S. R.; Golbon Haghighi, M.; Jamshidi, M.; Notash, B., Influence of the Diphosphine Coordination Mode on the Structural and Optical Properties of Cyclometalated Platinum (II) Complexes: An Experimental and Theoretical Study on Intramolecular Pt... Pt and π ... π Interactions. *Inorg. Chem.* **2018**, *57*, 5060-5073; (f) Nahaei, A.; Nabavizadeh, S. M.; Hosseini, F. N.; Hoseini, S. J.; Abu-Omar, M. M., Arene CH bond activation and methane formation by a methylplatinum (II) complex: Experimental and theoretical elucidation of mechanism. *New J. Chem.* **2019**, *43*, 8005-8014.
14. (a) Ojwach, S. O.; Ogwen, A. O.; Akerman, M. P., (Pyridyl) benzoazole palladium (II) complexes as homogeneous catalysts in hydrogenation of alkenes and alkynes. *Catal. Sci. Tech.* **2016**, *6*, 5069-5078; (b) Dey, T. K.; Ghosh, K.; Basu, P.; Molla, R. A.; Islam, S. M., Chloromethylated polystyrene immobilized ruthenium complex of 2-(2-pyridyl) benzimidazole catalyst for the synthesis of bioactive disubstituted ureas by carbonylation reaction. *New J. Chem.* **2018**, *42*, 9168-9176; (c) Guney, E.; Kaya, Y.; Yilmaz, V. T.; Gumus, S., Synthesis, experimental and theoretical characterization of palladium (II) and platinum (II) saccharinate complexes with 2-(2-pyridyl) benzimidazole. *Spect. Chim. Acta A: Mol. Biomol. Spect.* **2011**, *79*, 1171-1178; (d) Peng, D.; Zhang, W.; Tang, G.; Zhou, J.; Hu, J.; Xie, Q.; Zhong, C., Novel dye sensitizers of main chain polymeric metal complexes based on complexes of 2-(2'-pyridyl) benzimidazole derivative with Zn (II), Co (II): synthesis, characterization, and photovoltaic performance for dye-sensitized solar cells. *J. Iran. Chem. Soc.* **2015**, *12*, 397-404; (e) Serratrice, M.; Cinellu, M. A.; Maiore, L.; Pilo, M.; Zucca, A.; Gabbiani, C.; Guerri, A.; Landini, I.; Nobili, S.; Mini, E., Synthesis, structural characterization, solution behavior, and in vitro antiproliferative properties of a series of gold complexes with 2-(2'-pyridyl) benzimidazole as ligand: comparisons of gold (III) versus gold (I) and mononuclear versus binuclear derivatives. *Inorg. Chem.* **2012**, *51*, 3161-3171.
15. Jia, W. L.; McCormick, T.; Tao, Y.; Lu, J.-P.; Wang, S., New Phosphorescent Polynuclear Cu (I) Compounds Based on Linear and Star-Shaped 2-(2'-Pyridyl) benzimidazolyl Derivatives: Syntheses, Structures, Luminescence, and Electroluminescence. *Inorg. Chem.* **2005**, *44*, 5706-5712.

16. Zhong, C.; Guo, R.; Wu, Q., Design and syntheses of blue luminescent Cu (II) and Zn (II) polymeric complexes with 2-(2'-pyridyl) benzimidazole derivative ligand. *Reac. Func. Poly.* **2007**, *67*, 408-415.
17. Jia, W.-L.; Hu, Y.-F.; Gao, J.; Wang, S., Linear and star-shaped polynuclear Ru(II) complexes of 2-(2'-pyridyl) benzimidazolyl derivatives: syntheses, photophysical properties and red light-emitting devices. *Dalton Trans.* **2006**, 1721-1728.
18. Kamecka, A.; Kapturkiewicz, A.; Suwińska, K., Luminescent osmium (II) complexes with 2-(2-pyridyl)-benzimidazolate anion. *Inorg. Chem. Comm.* **2018**, *89*, 27-31.
19. (a) Yi, H.; Crayston, J. A.; Irvine, J. T., Ruthenium complexes of 2-(2'-pyridyl) benzimidazole as photosensitizers for dye-sensitized solar cells. *Dalton Trans.* **2003**, 685-691; (b) Wang, K.; Huang, L.; Gao, L.; Jin, L.; Huang, C., Synthesis, crystal structure, and photoelectric properties of Re (CO)₃ClL (L= 2-(1-ethylbenzimidazol-2-yl) pyridine). *Inorg. Chem.* **2002**, *41*, 3353-3358.
20. (a) Jamshidi, M.; Nabavizadeh, S. M.; Sepehrpour, H.; Hosseini, F. N.; Kia, R.; Rashidi, M., Cycloplatinated (II) complexes containing bridging bis (diphenylphosphino) acetylene: Photophysical study. *J. Lumin.* **2016**, *179*, 222-229; (b) Puddephatt, R., Compounds with platinum-metal bonds: A personal account. *J. Organomet. Chem.* **2017**, *849*, 268-278; (c) Jamali, S.; Nabavizadeh, S. M.; Rashidi, M., Binuclear cyclometalated organoplatinum complexes containing 1,1'-bis (diphenylphosphino) ferrocene as spacer ligand: kinetics and mechanism of Mel oxidative addition. *Inorg. Chem.* **2008**, *47*, 5441-5452; (d) Chen, J.-L.; Zeng, X.-H.; Ganesan, P.; He, L.-H.; Liao, J.-S.; Liu, S.-J.; Wen, H.-R.; Zhao, F.; Chi, Y., Heterobimetallic copper (I) complexes bearing both 1, 1'-bis (diphenylphosphino) ferrocene and functionalized 3-(2'-pyridyl)-1, 2, 4-triazole. *New J. Chem.* **2019**, *43*, 4261-4271; (e) Fallis, K. A.; Xu, C.; Anderson, G. K., Synthesis of unsymmetrical dpmm-bridged platinum and palladium A-frame complexes. *Organometallics* **1993**, *12*, 2243-2247.
21. (a) Shahsavari, H. R.; Aghakhanpour, R. B.; Babaghasabha, M.; Haghighi, M. G.; Nabavizadeh, S. M.; Notash, B., Photophysical properties of a series of cycloplatinated (II) complexes featuring allyldiphenylphosphane. *New J. Chem.* **2017**, *41*, 3798-3810; (b) Jamshidi, M.; Babaghasabha, M.; Shahsavari, H. R.; Nabavizadeh, S. M., The influence of thiolate ligands on the luminescence properties of cycloplatinated(II) complexes. *Dalton Trans.* **2017**, *46*, 15919-15927; (c) Aseman, M. D.; Nabavizadeh, S. M.; Niroomand Hosseini, F.; Wu, G.; Abu-Omar, M. M., Carbon–Oxygen Bond Forming Reductive Elimination from Cycloplatinated (IV) Complexes. *Organometallics* **2018**, *37*, 87-98; (d) Pazireh, S.; Aghakhanpour, R. B.; Fuertes, S.; Sicilia, V.; Hosseini, F. N.; Nabavizadeh, S. M., A double rollover cycloplatinated(II) skeleton: a versatile platform for tuning emission by chelating and non-chelating ancillary ligand systems. *Dalton Trans.* **2019**, *48*, 5713-5724; (e) Haghighi, M. G.; Rashidi, M.; Nabavizadeh, S. M.; Jamali, S.; Puddephatt, R. J., Cyclometalated organoplatinum (II) complexes: first example of a monodentate benzo [h] quinolyl ligand and a complex with bridging bis (diphenylphosphino) ethane. *Dalton Trans.* **2010**, *39*, 11396-11402; (f) Nabavizadeh, S. M.; Haghighi, M. G.; Esmaeilbeig, A. R.; Raoof, F.; Mandegani, Z.; Jamali, S.; Rashidi, M.; Puddephatt, R. J., Assembly of symmetrical or unsymmetrical cyclometalated organoplatinum complexes through a bridging diphosphine ligand. *Organometallics* **2010**, *29*, 4893-4899; (g) Fatemeh, N. H.; Farasat, Z.; Nabavizadeh, S. M.; Wu, G.; Abu-Omar, M. M., N-methylation versus oxidative addition using Mel in the reaction of organoplatinum (II) complexes containing pyrazine ligand. *J. Organomet. Chem.* **2019**, *880*, 232-240; (h) Niroomand Hosseini, F.; Nabavizadeh, S. M.; Abu-Omar, M. M., Which is the Stronger Nucleophile, Platinum or Nitrogen in Rollover Cycloplatinated (II) Complexes? *Inorg. Chem.* **2017**, *56*, 14706-14713; (i) Jamshidi, M.; Nabavizadeh, S. M.; Shahsavari, H. R.; Rashidi, M., Photophysical and DFT studies on cycloplatinated complexes: modification in luminescence properties by expanding of π -conjugated systems. *RSC Adv.* **2015**, *5*, 57581-57591.
22. Raoof, F.; Boostanizadeh, M.; Esmaeilbeig, A. R.; Nabavizadeh, S. M.; Aghakhanpour, R. B.; Ghiassi, K. B.; Olmstead, M. M.; Balch, A. L., Reactivity comparison of five- and six-membered cyclometalated platinum (II) complexes in oxidative addition reactions. *RSC Adv.* **2015**, *5*, 85111-85121.

23. Serratrice, M.; Maiore, L.; Zucca, A.; Stoccoro, S.; Landini, I.; Mini, E.; Massai, L.; Ferraro, G.; Merlino, A.; Messori, L., Cytotoxic properties of a new organometallic platinum (II) complex and its gold (I) heterobimetallic derivatives. *Dalton Trans.* **2016**, *45*, 579-590.
24. Romeo, R.; Monsù Scolaro, L.; Nastasi, N.; Mann, B. E.; Bruno, G.; Nicolò, F., trans-Chloromethylbis (dimethyl sulfoxide) platinum (II): X-ray Structure, Mechanism of Isomerization, and Its Use as a Precursor to Organoamine Complexes of Variable Geometrical Configurations. *Inorg. Chem.* **1996**, *35*, 7691-7698.
25. Sheldrick, G., *SADABS, Empirical Absorption Correction Program; University of Göttingen: Germany, 1997* **2005**.
26. SHELXTL PC, **Version 6.12, Bruker AXS Inc., Madison, WI, 2005**.
27. Frisch, M.; Trucks, G.; Schlegel, H.; Scuseria, G.; Robb, M.; Cheeseman, J.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G., *Gaussian 09, Revision A. 02; Gaussian, Inc: Wallingford, CT, 2009*
28. Hay, P. J.; Wadt, W. R., Ab initio effective core potentials for molecular calculations. Potentials for the transition metal atoms Sc to Hg. *J. Chem. Phys.* **1985**, *82*, 270-283.
29. Skripnikov, L., Chemissian: software to analyze spectra, build density maps and molecular orbitals. *Chemissian: software to analyze spectra, build density maps and molecular orbitals* **2016**, Version 4.
30. Nabavizadeh, S. M.; Niroomand Hosseini, F.; Nejabat, N.; Parsa, Z., Bismuth–Halide Oxidative Addition and Bismuth–Carbon Reductive Elimination in Platinum Complexes Containing Chelating Diphosphine Ligands. *Inorg. Chem.* **2013**, *52*, 13480-13489.
31. Falvello, L. R.; Fornies, J.; Gómez, J.; Lalinde, E.; Martín, A.; Martínez, F.; Moreno, M. T., Some platinum (II) complexes containing bis (diphenylphosphino) acetylene $\text{PPh}_2\text{C}\equiv\text{CPh}_2$: synthesis, characterisation and crystal structures. *J. Chem. Soc. Dalton Trans.* **2001**, 2132-2140.

SYNOPSIS TOC

An interesting cyclic tetranuclear Pt(II) complex, **2**, is synthesized by the reaction of **1** with $[\text{PtMe}(\text{Cl})(\text{DMSO})_2]$, followed by addition of dppac, bearing both the pbz (= 2-(2-pyridyl)benzimidazolate) and the dppac (= bis(diphenylphosphino)acetylene) ligands. The complex is the first example in which the pbz ligands act as both chelating and bridging to stabilize the tetraplatinum framework.

